1. For the next 24 hours:

- * Do not drive a car or do anything that requires alertness.
- * Do not stay by yourself or drink any alcohol.
- * Do not make important legal or business decisions.

Your alertness, memory and balance may be impaired today due to the sedation give for your procedure.

- 2. Notify your physician if you experience any of the following:
- - * Vomiting repeatedly or vomiting blood
- * Black tarry stools
- * Rectal bleeding of amount greater than 2 tablespoons
- * Redness, tenderness and swelling at IV site that lasts for more than 48 hours

* Shortness of breath * Rapid Heartrate.... Call 911 Chest pain

It is not uncommon after a Colonoscopy or after the first bowel movement to see a small amount of blood from the rectum or after an endoscopy or esophageal dilatation to see a small amount of blood in the sputum or to have a sore throat.

Please call you physician if you have problems. If you need a physician after hours, call the office and an answering service will help you locate the physician on call. If you are unable to reach a physician but feel your condition warrants a doctor's attention, go to an Emergency room near you.

3. Follow the instructions below:	
You may resume normal a	activities tomorrow.
You first meal should be	very light and non greasy (ex. soup and crackers).
Your second meal can	
be normal but avoid grea	sy, spicy or heavy foods. Regular diet day after
procedure.	
	w the instructions below for 5 7 10 14 days: vy lifting, climbing, pushing, pulling, contact
inflammatory drugs such as motrin,	es and no NSAID's (non-steroidal anti prosyn, etc.) You may take tylenol if needed for
	other tests, your results will be ready in 7 to 10 days sults
6. If your esophagus was dilated:	Clear liquids today. (No milk, or milk products, No jucies with pulp)
temperature)	Full liquids today. (Anything that is liquid at room Soft foods for days.

I, or other significant other, have be given and can verbalize these instructions.